

ISSN 1211-8788

Acta Musei Moraviae, Scientiae biologicae (Brno)
91: 69–82, 2006

The genus *Halictoxenos* Pierce, 1908 (Strepsiptera, Stylopidae) in the Czech Republic and Slovakia

JAKUB STRAKA¹, IGOR MALENOVSKÝ² & JAN BATELKA³

¹ Charles University in Prague, Department of Zoology, Viničná 7, CZ-128 44 Praha 2, Czech Republic;
e-mail: straka-jakub@vol.cz

² Moravian Museum, Department of Entomology, Hvězdoslavova 29a, CZ-627 00 Brno, Czech Republic;
e-mail: imalenovsky@mzm.cz

³ Nad Vodovodem 16, CZ-100 00 Praha 10, Czech Republic;
e-mail: janbat@centrum.cz

STRAKA J., MALENOVSKÝ I. & BATELKA J. 2006: The genus *Halictoxenos* Pierce, 1908 (Strepsiptera, Stylopidae) in the Czech Republic and Slovakia. *Acta Musei Moraviae, Scientiae biologicae* (Brno) 91: 69–82. – Based on a revision of museum material and recent collections, three species of the genus *Halictoxenos* Pierce, 1908 (Strepsiptera: Stylopidae) are recorded from the Czech Republic. They have been identified as *H. arnoldi* Perkins, 1918, *H. spencei* (Nassonov, 1893) and *H. tumulorum* Perkins, 1918. The latter two species were also found in Slovakia. Figures and diagnostic characters are provided for the females of the three species and information is given on their distribution and hosts. In both the countries, *H. arnoldi* has so far been found associated with *Lasioglossum* (*Lasioglossum*) *xanthopus*, *H. spencei* with eight solitary species from the “acarinate” lineage of *Lasioglossum*, subgenus *Evylaeus* and *H. tumulorum* with three species of *Halictus* (all Hymenoptera: Halictidae). *Lasioglossum* (*Evylaeus*) *sabulosum* (Warncke, 1986) and *L. (E.) sexstrigatum* (Schenck, 1870) are reported as new hosts for *H. spencei*. No morphological changes due to stylopization were observed on any of the examined hosts.

Key words. Strepsiptera, Stylopidae, *Halictoxenos*, parasitoids, Hymenoptera, Halictidae, *Lasioglossum*, *Halictus*, Central Europe, distribution, host associations.

Introduction

The Strepsiptera is a small order of insects consisting of entomophagous parasitoids. They are known to parasitize seven orders and thirty-five families of Insecta (KATHIRITHAMBY 1989). In Central Europe, twenty-one species of Strepsiptera have been found, of which fourteen species are associated with Hymenoptera (POHL & MELBER 1996).

The genus *Halictoxenos* Pierce, 1908 (Stylopidae) is confined to bees of the family Halictidae (the genera *Halictus* Curtis, 1833 and *Lasioglossum* Latreille, 1804) and is widely distributed in the Palaearctic, Nearctic, Afrotropical and Oriental Regions, and Australia (HOFENEDER & FULMEK 1943, KIFUNE 1982, POHL & KINZELBACH 1995, KATHIRITHAMBY & TAYLOR 2005). The European species of *Halictoxenos* were last addressed by KINZELBACH (1978), who summarized information on their hosts, distribution and synonymy, and provided redescriptions and keys for determination. He recognized three species in Europe: *H. arnoldi* Perkins, *H. spencei* Nassonov and *H. tumulorum* Perkins.

J. STRAKA, I. MALENOVSKÝ & J. BATELKA

Data on the Czech and Slovak Strepsiptera, including *Halictoxenos*, have been scarce. OGLOBLIN (1925), while describing the new species *Halictoxenos nitidiusculus* Ogloblin from Poland, also listed a record of two females from former Czechoslovakia (Bohemia). HOFENEDER & FULMEK (1943) cited *Halictoxenos tumulorum*, parasitic on *Halictus tumulorum*, from Kolín (Bohemia). GÜNTHER (1947) received stylized material from several Czech hymenopterologists and compiled a list of hosts of Strepsiptera in Czechoslovakia that included eight different species of *Halictus* and *Lasioglossum*. He identified the corresponding Strepsiptera as *Halictoxenos nitidiusculus*, *H. spencei*, *H. tumulorum* and *H. sp.* and later reported the first three taxa in the checklist of the Czechoslovak insect fauna (GÜNTHER 1977). Finally, based on information on the hosts but without directly examining the specimens KINZELBACH (1978) associated each of these previously published records with *Halictoxenos arnoldi*, *H. spencei* or *H. tumulorum*.

Recently, we checked the Günther collection of Strepsiptera deposited in the National Museum, Prague. We also found some other unpublished *Halictoxenos* material in this institution and in the Moravian Museum Brno and collected new material in the field. We report all available information on the Czech and Slovak *Halictoxenos* in this paper.


Material and methods

In the material section, localities are supplemented by a code number (in parentheses), which refers to the map field of the Central-European grid for mapping flora and fauna (EHRENDORFER & HAMANN 1965, adapted by NOVÁK 1989 and PRUNER & MÍKA 1996). The nomenclature of the hosts (Hymenoptera: Halictidae) is used according to SCHWARZ *et al.* (1996). Jakub Straka identified all the hosts.

For Strepsiptera, we employ the nomenclature and synonymy used by KINZELBACH (1978). The morphological terminology is according to KATHIRITHAMBY (1991). Measurements are given in mm and were made from slide-mounted specimens. Slides were photographed using a digital camera mounted on a microscope, scaled and measured by image analysis (Olympus QuickPHOTO PRO).

The Strepsiptera material is mounted on slides (mostly in Canada Balsam or glycerine-gelatine), conserved dry in the hosts or in ethanol. It is deposited (mostly together with the hosts) in the following collections:

JB	Jan Batelka private collection, Prague
JS	Jakub Straka private collection, Prague
MMB	Moravian Museum, Brno (coll. Strepsiptera)
NMP	National Museum, Prague (coll. Günther and general collection)



Halictoxenos (Strepsiptera, Stylopidae) in the Czech Republic and Slovakia






Fig. 1. *Halictoxenos* spp., female cephalothorax. A – *H. arnoldi*, Bohemia: Tábor, from *Lasioglossum xanthopus*. B – *H. arnoldi*, Bohemia: Chýnice, from *Lasioglossum xanthopus*. C – *H. tumulorum*, Slovakia: Kamenica nad Hronom, from *Halictus kessleri*. D – *H. tumulorum*, Slovakia: Štúrovo, from *Halictus simplex*.





J. STRAKA, I. MALENOVSKÝ & J. BATELKA






Fig. 2. *Halictoxenos spencei*, female cephalothorax. A – Slovakia: Dražovce, from *Lasioglossum limbellum*. B – Moravia: Čejč, from *Lasioglossum nitidiusculum*. C – Moravia: Hradec nad Moravicí, from *Lasioglossum parvulum*. D – Moravia: Bzenec, from *Lasioglossum sexstrigatum*.





Halictoxenos (Strepsiptera, Stylopidae) in the Czech Republic and Slovakia




Fig. 3. *Halictus simplex* parasitized by *Halictoxenos tumulorum* (Slovakia: Štúrovo). Arrows point at the female cephalothorax extruding exterior of host. A – dorsal view of the host. B – detail of the host abdomen, in dorso-lateral view. C – detail of the cephalothorax of *H. tumulorum*.

Results


Halictoxenos arnoldi Perkins, 1918 (Figs 1A, B; 4A)

Material examined. Czech Republic, Bohemia: Chýnice (6051), 17.iv. (year unknown, second half of the 20th century), 1 ♀, in 1 ♀ *L. xanthopus*, B. Tkalcù lgt. (NMP, host dry-mounted, parasitoid on slide); Tábor (6553–54), ix.1939, 2 ♀♀, in 1 ♂ *Lasioglossum xanthopus*, A. Hoffer lgt. (NMP, host dry-mounted, parasitoids on slides).

Published data. GÜNTHER (1947): 72 (Tábor, as *Halictoxenos* sp. and *H. nitidiusculus*); KINZELBACH (1978): 106, 108 (Tábor, as *H. arnoldi* and *H. spencei*).

Hosts. *Lasioglossum (Lasioglossum) xanthopus* (Kirby, 1802). According to KINZELBACH (1978) probably also associated with other *Lasioglossum (Lasioglossum)* spp.: *L. costulatum* (Kriechbaumer, 1873), *L. leucozonium* (Schrank, 1781) and *L. quadrinotatum* (Kirby, 1802).

Distribution. Czech Republic, Germany, Great Britain, Hungary and Turkey (KINZELBACH 1978, POHL 2004). In the Czech Republic so far known only from Central and South Bohemia (Fig. 4A), not yet recorded from Slovakia.



J. STRAKA, I. MALENOVSKÝ & J. BATELKA




Fig. 4. Distribution of *Halictoxenos arnoldi* (black triangles) and *H. tumulorum* (black circles). A – Czech Republic. B – Slovakia.

Halictoxenos (Strepsiptera, Stylopidae) in the Czech Republic and Slovakia*Halictoxenos spencei* (Nassonov, 1893)

(Figs 2A–D; 5A, B)

Nomenclatural note: The original spelling of the specific name is “*spencii*”. The subsequent spelling “*spencei*” introduced by KINZELBACH (1969) may be preserved in accordance with ICBN (1999), Art. 33.3.1.

Synonymy: *Halictoxenos calceati* Noskiewicz & Poluszyński, 1925; *Halictoxenos cylindrici* Perkins, 1918; *Halictoxenos (Halictostylops) nitidiusculus* Ogleblin, 1925; *Halictoxenos puncticollis* Noskiewicz & Poluszyński, 1925. See HOFENEDER & FULMEK (1943) and KINZELBACH (1978) for further notes.

Material examined. Czech Republic, Bohemia: Bynovec (5151), 28.vi.2005, 1 ♀, in 1 ♀ *Lasioglossum punctatissimum*, coloured pan traps, L. Blažej lgt. (JS, host dry-mounted, parasitoid specimen lost during preparation); Slaný (5750), 15.vi.1923, 1 ♀, in 1 ♀ *Lasioglossum nitidiusculum*, S. Hrabě lgt. (NMP, dry-mounted in host); Praha-Dolní Liboc, Divoká Šárka Nature Reserve (5951), 9.iv.1998, 1 ♀, in 1 ♀ *Lasioglossum parvulum*, M. Kolíško lgt. (JB, dry-mounted in host); Praha-Dubeček (5953), 5.vi.2005, 2 ♀♀, in 1 ♀ *Lasioglossum sabulosum*, observed and collected on *Salvia officinalis* flowers in a garden, in late afternoon at 4 p.m. in cold weather (only *Apis mellifera* Linnaeus, 1758 and *Bombus* sp. were active in the same time), J. Batelka lgt. (JS, in ethanol); Praha-Radotín (6052), 20.iv.1941, 1 ♀, in 1 ♀ *Lasioglossum parvulum*, Píša lgt. (NMP, dry-mounted in host); Zvole, Zvolská homole (6052), 13.v.2006, 18 ♀♀, in 13 ♀♀ *Lasioglossum parvulum* and 2 ♀♀ *Lasioglossum pygmaeum* (1–3 Strepsiptera specimens per host), in hosts’ nesting area, in cloudy cold weather, on the ground or in flight, J. Batelka & J. Straka lgt. (JS, in ethanol); Strančice (6054), vi.1949, 3 ♀♀, in 1 ♀ *Lasioglossum parvulum*, V. Kočmíd lgt. (NMP, dry-mounted in host); Písek (6650), 14.vi.1930, 1 ♀, in 1 ♀ *Lasioglossum nitidiusculum*, K. Táborský lgt. (NMP, dry-mounted in host).

Czech Republic, Moravia: Hradec (Silesia) [= Hradec nad Moravici] (6173), 30.iv.1933, 2 ♀♀, in 1 ♀ *Lasioglossum parvulum*, J. Palásek lgt. (NMP, host dry-mounted, parasitoids on slides); Tišnov (6664), 21.iv.1936, 3 ♀♀, in 2 ♀♀ *Lasioglossum parvulum*, A. Hoffer lgt. (NMP, dry-mounted in host); Brno-Bystrc (6765), 5.v.1943, 2 ♀♀, in 1 ♀ *Lasioglossum nitidiusculum*, J. Šnoflák lgt. (MMB, dry-mounted in host); Velehrad (6870), 5.viii.1940, 1 ♀, in 1 ♀ *Lasioglossum villosulum*, V. Zavadil lgt. (NMP, host dry-mounted, parasitoid on slide); Pouzdřany (7065), 8.vii.1938, 2 ♀♀, in 1 ♂ *Lasioglossum nitidiusculum*, V. Zavadil lgt. (NMP, dry-mounted in host); ibid., 13.vii.2005, 1 ♀, in 1 ♀ *Lasioglossum villosulum*, in a vineyard, collected running on the ground, in cold and humid weather at 6 p.m.; P. Bogush & J. Straka lgt. (JS, in ethanol); ibid., 24.iv.2006, 1 ♀, in 1 ♀ *Lasioglossum punctatissimum*, coloured pan traps, J. Batelka & J. Straka lgt. (JS, in ethanol); Čejč u Hodonína (7067), 7.ix.1941, 1 ♀, in 1 ♀ *Lasioglossum nitidiusculum*, V. Zavadil lgt. (NMP, host dry-mounted, parasitoid on slide); ibid., v.1941, 3 ♀♀, in 2 ♀♀ *Lasioglossum nitidiusculum*, collector unknown (NMP, hosts dry-mounted, parasitoids on slides); Horovany (7067–68), 29.vi.1944, 2 ♀♀ and first instar larvae, in 1 ♀ *Lasioglossum nitidiusculum*, O. Šustera lgt. (NMP, host dry-mounted, parasitoids on slides); Bzenec (7069), 6.vii.1942, 2 ♀♀, in 1 ♀ *Lasioglossum sexstrigatum*, O. Šustera lgt. (NMP, host dry-mounted, parasitoids on slides); Dolní Dunajovice (7165), 6.v.2006, 1 ♀, in 1 ♀ *Lasioglossum nitidiusculum*, swept from *Stellaria media* in a vineyard, at 1 p.m., in sunny weather, J. Straka lgt. (JS, in ethanol); Dolní Věstonice (7165), 12.vi.2006, 1 ♀, in 1 ♀ *Lasioglossum pygmaeum*, on *Bryonia alba* flower, at 10 a.m., in sunny weather, J. Straka lgt. (JS, in ethanol).

Slovakia: Dražovce near Nitra (7674), 25.viii.1948, 1 ♀, in 1 ♀ *Lasioglossum limbellum*, O. Šustera lgt. (NMP, host dry-mounted, parasitoid on slide); Parkan [= Štúrovo] (81–8278), 7.vii.1946, in 1 ♂ *Lasioglossum nitidiusculum*, O. Šustera lgt. (NMP, dissected host preserved dry, parasitoid not found)..

Published data. OGLOBLIN (1925): 116 (Slaný, as *H. nitidiusculus*); GÜNTHER (1947): 72 (Horovany, Velehrad, as *H. spencii*; Čejč, as *H. nitidiusculus*; Parkán, Radotín, as *Halictoxenos* sp.), KINZELBACH (1978): 108–109 (Slaný, Čejč, Horovany, Velehrad; Písek, Pouzdřany – the latter two localities misspelled as “Přseg” and “Przdwiány”, as *H. spencii*; Parkán, Radotín, as *H. spencii* or *H. tumulorum*). Based on our examination of the Günther specimens and their labels, the record of *H. spencii* from *L. nitidiusculum*, Bohemia: Tábor listed by GÜNTHER (1947) is a mistake and refers to *H. arnoldi* and *Lasioglossum (L.) xanthopus*, respectively. The hosts of the *Halictoxenos spencii* specimens from Velehrad published by GÜNTHER (1947) as *Halictus minutus* [= *Lasioglossum parvulum*] and from Parkán as *Halictus convexiusculus* [= *Lasioglossum convexiusculus* (Schenck, 1853)] were both misidentified and belong to *Lasioglossum villosulum* and *Lasioglossum nitidiusculum*, respectively. The specimens of *Halictus morio* [= *Lasioglossum morio* (Fabricius, 1793)] and *Halictus viridiaeneus* [= *Lasioglossum aeratum* (Kirby, 1802)] from Radotín which GÜNTHER (1947) and

J. STRAKA, I. MALENOVSKÝ & J. BATELKA




Fig. 5. Distribution of *Halictoxenos spencei* (black circles). A – Czech Republic. B – Slovakia.

Halictoxenos (Strepsiptera, Stylopidae) in the Czech Republic and Slovakia

KINZELBACH (1978) had referred to were also found in the Günther collection in NMP and checked by us. However, we failed to find any traces of their stylopization.

Hosts. Restricted to the subgenus *Evylaeus* Robertson, 1902 of the genus *Lasioglossum*. In the Czech Republic and Slovakia so far recorded from *Lasioglossum* (*Evylaeus*) *limbellum* Morawitz, 1876; *L. (E.) nitidiusculum* (Kirby, 1802); *L. (E.) parvulum* (Schenck, 1853); *L. (E.) punctatissimum* (Schenck, 1859); *L. (E.) pygmaeum* (Schenck, 1853); *L. (E.) sabulosum* (Warncke, 1986); *L. (E.) sexstrigatum* (Schenck, 1870); *L. (E.) villosulum* (Kirby, 1802). *L. (E.) sabulosum* and *L. (E.) sexstrigatum* are reported as the hosts of *Halictoxenos spencei* for the first time. For a full list of confirmed and unconfirmed hosts see KINZELBACH (1978).

Distribution. Widely distributed in the western Palaearctic region. Reported from Austria, Belgium, Great Britain, Canary Islands, Czech Republic, Denmark, Finland, France, Germany, Greece (mainland and Crete), Hungary, Ireland, Italy (including Sardinia), Norway, Poland, Slovakia, Spain, Ukraine, and Turkey (KINZELBACH 1978, POHL 2004, BATELKA & STRAKA 2005, RONAYNE & O'CONNOR 2006). Distribution in the Czech Republic and Slovakia is shown in Fig. 5A, B.

Halictoxenos tumulorum Perkins, 1918 (Figs 1C, D; 3; Fig. 4A, B)

Synonymy: *Halictoxenos rubicundi* Noskiewicz & Poluszyński, 1924; *Halictoxenos sajoi* Noskiewicz & Poluszyński, 1924; *Halictoxenos simplicis* Noskiewicz & Poluszyński, 1935; *Halictoxenos ulrichi* Hofeneder, 1939.

Material examined. Czech Republic, Bohemia: Praha-Stodůlky (5951), 21.viii.–9.ix.2005, 1 ♀, in 1 ♂ *Halictus tumulorum*, coloured pan trap, J. Straka lgt. (JS, in ethanol). Praha-Dubeček (5953), 18.iv.–25.iv.2006, 1 ♀, in 1 ♂ *Halictus tumulorum*, coloured pan trap, J. Batelka lgt. (JS, in ethanol); ibid., 25.iv.–4.v.2006, 2 ♀♀, in 1 ♂ *Halictus tumulorum*, coloured pan traps, J. Batelka lgt. (JS, in ethanol); ibid., 7.viii.–14.viii.2006, 1 ♀, in 1 ♂ *Halictus tumulorum*, coloured pan trap, J. Batelka lgt. (JS, in ethanol).

Czech Republic, Moravia: Tasovice (7162), 24.vi.2006, 18 ♀♀, in 14 ♀♀ *Halictus kessleri* (1–3 Strepsiptera specimens per host), swept from *Berteroia incana*, J. Batelka & J. Straka lgt. (JS, in ethanol); Jaroslavice (7263), 28.iv.2006, 3 ♀♀, in 2 ♀♀ *Halictus kessleri*, swept from *Erysimum* sp., J. Straka lgt. (JS, in ethanol).

Slovakia: Kamenica nad Hronom (8178), 22.vi.1946, 2 ♀♀, in *Halictus kessleri*, Šusterá lgt. (NMP, host dry-mounted, parasitoids on slides); Štúrovo, soutok Hronu s Dunajem (8178), 27.vii.1974, 1 ♀, in 1 ♂ *Halictus simplex*, B. Tkalcú leg. (JS, host dry-mounted, parasitoid on slide).

Published data. HOFENEDER & FULMEK (1943): 34 (Kolín, as *H. tumulorum*); GÜNTHER (1947): 72 (Kolín, as *H. tumulorum*, Kamenica nad Hronom, as *H. sp.*), KINZELBACH (1978): 106, 108 (Kolín, Morawa, Kamenica nad Hronom, as *H. tumulorum*). The host of the *Halictoxenos* specimen from Kamenica nad Hronom published by GÜNTHER (1947) as *Halictus subauratus* (Rossi, 1792) was misidentified and belongs to *H. kessleri*. We have not found the specimens of *H. tumulorum* from Kolín (HOFENEDER & FULMEK 1943, GÜNTHER 1947) and "Morawa" [= Moravia?] (Günther in litt., KINZELBACH 1978) in NMP; these records thus remain unchecked.

Hosts. In the Czech Republic and Slovakia found to date only in *Halictus* (*Seladonia*) *kessleri* Bramson, 1879, *H. (S.) tumulorum* (Linnaeus, 1758) and *H. (Halictus) simplex* Blüthgen, 1923. According to KINZELBACH (1978), parasitizes a wide range of species from the subgenera *Halictus* and *Seladonia* Robertson, 1918 of the genus *Halictus* and perhaps also some species of *Lasioglossum* (*Evylaeus*).

J. STRAKA, I. MALENOVSKÝ & J. BATELKA

Distribution. Widely distributed in the western Palaearctic region. Reported from Austria, Great Britain, Canary Islands, Czech Republic, Finland, France, Germany, Hungary, Ireland, Italy, Netherlands, Portugal, Russia (NW European part), Slovakia, Spain, Ukraine, Turkey, and North Africa (KINZELBACH 1978, POHL 2004, SMIT & SMIT 2005, RONAYNE & O'CONNOR 2006). In the Czech Republic so far found in the warm regions of Central Bohemia and South Moravia. It has also been recorded from southern Slovakia (Fig. 4A, B).

Discussion

A taxonomic study of the European *Halictoxenos* is difficult because males are scarce. Facultative parthenogenesis has been mentioned as a possible means of reproduction (KINZELBACH 1978). Males have been described so far only in *H. spencei* and *H. tumulorum* (including *H. ulrichi*, a synonym of *H. tumulorum*; for the descriptions of males see KINZELBACH 1978). The descriptions of *H. arnoldi*, as well as several other taxa that are currently treated as synonyms, are based only on females (PERKINS 1918a, b, OGLOBLIN 1925, NOSKIEWICZ & POLUSZYŃSKI 1924, 1935). Furthermore, all the specimens examined from the Czech Republic and Slovakia are females. Females of Strepsitera, suborder Stylopida, are neotenic and being devoid of many adult features (e.g. wings, antennae, legs and external genitalia) they are often difficult to identify down to species. Moreover, their size varies according to the size of the host (KATHIRITHAMBY 1989). The measurements of our specimens indeed did not yield any help for identification purposes since the morphometric characters largely overlap in all the three species (Tables 1–2). The outline of the cephalothorax is more or less constant within a species, even if it is also subject to a certain variation (Figs 1–2). *H. tumulorum* can be differentiated from *H. arnoldi* and *H. spencei* by the regularly arcuate brood passage opening and lack of pale to translucent spots medially on the cephalothorax forward of an imaginary line connecting the spiracles. These spots correspond to the openings of Nassonov's pheromone glands (KINZELBACH 1978). In *H. tumulorum* the Nassonov glands are distinct as a pair of pale oblong fields at the cephalothorax base on each side of the midline, behind the spiracles. In contrast, both *H. arnoldi* and *H. spencei* share a slightly angular brood passage opening and several (ca. 9–15) small irregular Nassonov gland spots in the middle of cephalothorax forward of the line connecting the spiracles. These spots are translucent and very distinct on all specimens of *H. spencei* examined, whereas they are more difficult to observe in *H. arnoldi*. Contrary to KINZELBACH (1978) and in agreement with the original description (PERKINS 1918b), we were able to locate a series of spots (slightly paler than the rest of cephalothorax) in the middle of cephalothorax in two out of the three *Halictoxenos* specimens from *Lasioglossum xanthopus* studied. Morphological characters be useful for the identification of *H. arnoldi*, *H. spencei* and *H. tumulorum* are summarized in Table 3. A more detailed study of the systematics of the genus (including relationships to non-European species), based only on female morphology, is probably impossible due to the

Halictoxenos (Strepsiptera, Stylopidae) in the Czech Republic and Slovakia

Species	N	CL	CW	MW	BW	AW
<i>arnoldi</i>	3	0.54–0.88	0.57–0.85	0.08–0.09	0.21–0.25	0.38–0.57
<i>spencsei</i>	12	0.47–0.69	0.44–0.70	0.07–0.12	0.17–0.31	0.28–0.58
<i>tumulorum</i>	3	0.57–0.82	0.51–0.83	0.08–0.14	0.18–0.30	0.36–0.57


Table 1. Measurements of *Halictoxenos* spp., females. N – number of measured specimens. CL – cephalothorax length (from cephalothorax apex to abdomen base). CW – cephalothorax maximum width. MW – distance between the mandibles. BW – brood passage opening width. AW – abdomen base width.

Species	N	CL/CW	BW/CW	AW/CW
<i>arnoldi</i>	3	0.93–1.04	0.29–0.38	0.66–0.74
<i>spencsei</i>	12	0.99–1.29	0.38–0.45	0.58–0.94
<i>tumulorum</i>	3	0.98–1.14	0.36–0.40	0.68–0.72

Table 2. Ratios of *Halictoxenos* spp., females. See the key to Table 1 for explanations.

Species	Cephalothorax shape	Brood passage opening	Pale spots on cephalothorax (Nassonov's gland fields)	Hosts
<i>arnoldi</i>	posterior edges ± sharply angular	slightly angular	numerous hardly distinct small spots situated medially and forward of an imaginary line connecting the spiracles	<i>Lasioglossum xanthopus</i> ; ?other <i>Lasioglossum</i> (<i>Evylaeus</i>) spp.
<i>spencsei</i>	posterior edges ± obtusely angular	slightly angular	numerous clearly distinct small spots situated medially and forward of an imaginary line connecting the spiracles	<i>Lasioglossum</i> (<i>Evylaeus</i>) spp.
<i>tumulorum</i>	posterior edges ± narrowly rounded	regularly arcuate	two large oblong spots situated on each side of the midline at the cephalothorax base behind an imaginary line connecting the spiracles	<i>Halictus</i> spp.; ? <i>Lasioglossum</i> (<i>Evylaeus</i>) spp.

Table 3. Diagnostic characters of females of European *Halictoxenos* spp.



J. STRAKA, I. MALENOVSKÝ & J. BATELKA


lack of other observable characters. A study of molecular characters would certainly be one of the best approaches to this problem.


The records from the Czech Republic and Slovakia would suggest that *Halictoxenos tumulorum* is restricted to the genus *Halictus*. KINZELBACH (1978) listed also some species of *Lasioglossum (Evylaeus)* as being hosts of *H. tumulorum*, among them also *Lasioglossum (Evylaeus) limbellum*. The female Strepsiptera specimen from *L. (E.) limbellum* we examined from Slovakia, however, clearly corresponds to *Halictoxenos spencei* (Fig. 2A). *L. (E.) limbellum* as well as all the remaining Czech and Slovak host bees of *H. spencei* belong to the “acarinate” lineage within the subgenus *Evylaeus*, which is treated as a secondary solitary group by DANFORTH *et al.* (2003). Besides the “acarinate” species, several species from the “carinate” group of *Evylaeus* [e.g. *L. calceatum* (Scopoli, 1763), *L. fulvicorne* (Kirby, 1802) and *L. pauxillum* (Schenck, 1853)] can also be parasitized by *H. spencei* (PERKINS 1918a, NOSKIEWICZ & POLUSZYŃSKI 1924, KINZELBACH 1978). These “carinate” species are primarily eusocial but each of them may be polymorphic in degree of social behaviour, up to solitariness (RICHARDS 2000). The only known host of *Halictoxenos arnoldi* in the Czech Republic has so far been *Lasioglossum (Lasioglossum) xanthopus*. This species seems to be the main host of *H. arnoldi* in Germany as well (SAURE 2003).

Specimens of Hymenoptera parasitized by Strepsiptera often show morphological changes due to the effects of the parasitoid on the host. In some characters (e.g. face maculation, antenna segmentation, form and length of hairs on legs and abdomen) stylopized males tend to resemble normal females and stylopized females resemble normal males. These morphological modifications and “intersexual” specimens have been well-documented in stylopized bees from the genus *Andrena* Fabricius, 1775 and some solitary vespids and digger wasps (PÉREZ 1886, PIERCE 1909, SALT 1927, 1931, KATHIRITHAMBY 1989). In contrast, we observed no such modifications of secondary sexual characters on any of the host specimens of *Halictus* and *Lasioglossum* parasitized by *Halictoxenos* and examined by us.

Acknowledgements


We are grateful to Jiří Hájek and Jan Macek (National Museum, Prague) for providing us access to the NMP museum collection, Luboš Dembický (Moravian Museum, Brno) for the macrophotographs of *Halictus simplex*, Petr Kment (National Museum, Prague) for help with literature and Vítězslav Kubáň (Moravian Museum, Brno) for valuable comment. Our work was partly supported by grant 0021620828 from the Ministry of Education, Youth and Sports of the Czech Republic to the Faculty of Science, Charles University, Prague and grant MK00009486201 from the Ministry of Culture of the Czech Republic to the Moravian Museum, Brno.






References

- BATELKA J. & STRAKA J. 2005: Several records of Xenidae and Stylopidae from the West Palaearctic region (Strepsiptera). *Bulletin de la Société entomologique de France* **110**: 403–406.
- DANFORTH B. N., CONWAY L. & JI S. 2003: Phylogeny of eusocial Lasiglossum reveals multiple losses of eusociality within a primitively eusocial clade of bees (Hymenoptera: Halictidae). *Systematic Biology* **52**: 23–36.
- EHRENDORFER F. & HAMANN U. 1965: Vorschläge zu einer floristischen Kartierung von Mitteleuropa. *Berichte der Deutschen botanischen Gesellschaft* **78**: 35–50.
- GÜNTHER V. 1947: Předběžná zpráva o řádu Strepsiptera v Československu. Preliminary report on the order Strepsiptera in Czechoslovakia. *Acta entomologica Musei nationalis Pragae* **25**: 69–72 (in Czech and English).
- GÜNTHER V. 1977: Strepsiptera. P. 123. In: Dlabola J. (ed.): *Enumeratio insectorum bohemoslovakiæ*. Check List tschechoslowakische Insektenfauna. *Acta faunistica entomologica Musei nationalis Pragae* **15, Suppl. 4**: 1–158.
- HOFENEDER K. & FULMEK L. 1943: Verzeichnis der Strepsiptera und ihrer Wirte. *Arbeiten über physiologische und angewandte Entomologie* **10**: 32–58.
- ICZN 1999: International code of zoological nomenclature. Fourth Edition. The International Trust for Zoological Nomenclature, London, 271 pp.
- KATHIRITHAMBY J. 1989: Review of the order Strepsiptera. *Systematic Entomology* **14**: 41–92.
- KATHIRITHAMBY J. 1991: Strepsiptera. Pp. 684–695. In: Naumann I. D. (ed.): *Insects of Australia*. Second edition, Melbourne University Press, 1137 pp.
- KATHIRITHAMBY J. & TAYLOR S. J. 2005: A new species of Halictophagus (Insecta: Strepsiptera: Halictophagidae) from Texas, and a checklist of Strepsiptera from the United States and Canada. *Zootaxa* **1056**: 1–18.
- KIFUNE T. 1982: Four new species of the genus Halictoxenos Pierce from Japan (Strepsiptera, Stylopidae). *Esakia* **19**: 151–180.
- KINZELBACH R. K. 1969: Familie: Stylopidae, Fächerflügler. Pp. 139–159. In: Freude H., Harde K. W. & Lohse G. A.: *Die Käfer Mitteleuropas*, Band 8 (Teredilia, Heteromera, Lamellicornia). Goecke & Evers, Krefeld, 388 pp.
- KINZELBACH R. K. 1978: Insecta. Fächerflügler (Strepsiptera). Die Tierwelt Deutschlands 65. VEB Gustav Fischer Verlag, Jena, 166 pp.
- NOVÁK I. 1989: Seznam lokalit a jejich kódů pro síťové mapování entomofauny Československa [List of localities and their codes for entomofaunistic grid mapping of Czechoslovakia]. *Zprávy Československé společnosti entomologické při ČSAV* **25**: 3–84 (in Czech).
- NOSKIEWISZ J. & POLUSCZYŃSKI G. 1925: Neue Arten der Strepsipterengattung Halictoxenos Pierce. *Polskie pismo entomologiczne* **3 (1924)**: 182–188.
- NOSKIEWISZ J. & POLUSCZYŃSKI G. 1935: Embryologische Untersuchungen an Strepsiteren. II. Teil. Polymobryonie. *Zoologica Poloniae* **1**: 53–94.
- OGŁOBLIN A. 1925: New and little known Strepsiptera from Poland. *Polskie pismo entomologiczne* **3 (1924)**: 113–122.
- PERKINS R. C. L. 1918a: Synopsis of British Strepsiptera of the genera Stylops and Halictoxenos. *Entomologist's Monthly Magazine* **54**: 67–76.
- PERKINS R. C. L. 1918b: Halictoxenos arnoldi, an undescribed British Stylopid. *Entomologist's Monthly Magazine* **54**: 107–108.
- PÉREZ J. 1886: Des effets du parasitisme des Stylops sur les apiaries du genre Andrena. *Actes de la Société linnéenne de Bordeaux* **40**: 21–60.
- PIERCE W. D. 1909: A monographic revision of the twisted winged insects comprising the order Strepsiptera. *Bulletin of the United States National Museum* **66**: xii, 1–232, 15 pl.





J. STRAKA, I. MALENOVSKÝ & J. BATELKA

- POHL H. 2004: Fauna Europaea: Strepsiptera. Fauna Europaea, version 1.1, <http://www.faunaeur.org> (accessed 12 September 2006).
- POHL H. & KINZELBACH R. 1995: A new species of Halictoxenos Pierce (Strepsiptera: Stylopida: Stylopidae) from South Africa. *African Entomology* **3**: 73–76.
- POHL H. & MELBER A. 1996: Verzeichnis der mitteleuropäischen Fächerflügler und die Beschreibung einer neuen Art der Gattung Malayaxenos Kifune 1981 (Insecta: Strepsiptera). *Senckenbergiana biologica* **75**: 171–180.
- PRUNER L. & MÍKA P. 1996: Seznam obcí a jejich částí v České republice s čísly mapových polí pro síťové mapování fauny. [List of settlements in the Czech Republic with associated map field codes for faunistic grid mapping system]. *Klapalekiana* **32** (Suppl.): 1–115 (in Czech, English summary).
- RICHARDS M. H. 2000: Evidence for geographic variation in colony social organization in an obligately social sweat bee, *Lasioglossum malachurum* Kirby (Hymenoptera; Halictidae). *Canadian Journal of Zoology* **78**: 1259–1266.
- RONAYNE C. & O'CONNOR J. P. 2006: *Halictoxenos tumulorum* Perkins & *H. spencei* Nassonov (Strepsipt., Stylopidae) new to Ireland. *Entomologist's Monthly Magazine* **142**: 92.
- SALT G. 1927: The effects of stylopization on aculeate Hymenoptera. *Journal of Experimental Zoology* **48**: 223–331.
- SALT G. 1931: A further study of the effects of stylopization on wasps. *Journal of Experimental Zoology* **59**: 133–166.
- SAURE C. W. 2003: Insekten am konventionellen und transgenen Raps Blütenbesuch, Pollentransfer und Auskreuzung. Insects in transgenic and conventional oilseed rape. PhD thesis, Freie Universität Berlin, Fachbereich Biologie, 158 pp. (available on-line on <http://www.diss.fu-berlin.de/2003/236/index.html>).
- SCHWARZ M., GUSENLEITNER F., WESTRICH P. & DATHE H. H. 1996: Katalog der Bienen Österreichs, Deutschlands und der Schweiz (Hymenoptera, Apidae). *Entomofauna*, Suppl. **8**: 1–398.
- SMIT J. T. & SMIT J. 2005: De waaiervleugeligen (Strepsiptera) van Nederland. *Entomologische Berichten* **65**: 43–51.

